Table 18. Lunch¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic

Race/ethnicity and age	Percent reporting ³	Energy	Protein	2		ate sugars fiber		Saturated fat	Mono- unsaturated fat	Poly- unsaturated fat	
(years)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	
Non-Hispanic White:		1									
2 - 5	96* (1.4) 94 (1.6) 85 (2.5) 80 (1.7)	24 (1.0) 25 (0.8) 25 (0.9) 23 (0.5)	26 (1.0) 28 (1.1) 27 (1.0) 26 (0.5)	21 (0.9) 23 (0.9) 23 (0.9) 22 (0.5)	18 (0.8) 21 (1.2) 20 (1.1) 17 (0.7)	24 (1.3) 25 (1.0) 24 (1.0) 24 (0.7)	27 (1.5) 27 (1.1) 26 (1.0) 25 (0.5)	25 (1.0) 26 (1.2) 25 (1.2) 24 (0.6)	28 (1.8) 26 (1.2) 26 (1.2) 25 (0.5)	29 (2.3) 29 (1.3) 28 (1.1) 27 (0.6)	
2 and over	82 (1.5)	23 (0.4)	26 (0.5)	22 (0.5)	18 (0.6)	24 (0.6)	25 (0.5)	24 (0.6)	25 (0.5)	27 (0.5)	
Non-Hispanic Black:											
2 - 5	93* (1.7) 83 (2.7) 73 (2.6) 68 (1.4)	28 (1.9) 23 (1.3) 26 (1.5) 23 (0.5)	30 (1.7) 27 (1.2) 28 (1.6) 26 (0.7)	26 (2.1) 23 (1.5) 24 (1.4) 22 (0.7)	22 (2.5) 22 (1.8) 22 (1.3) 19 (0.8)	28 (2.5) 27 (2.0) 28 (1.7) 24 (0.7)	30 (2.0) 23 (1.3) 26 (1.7) 25 (0.5)	30 (2.0) 22 (1.4) 26 (1.7) 24 (0.5)	31 (2.1) 23 (1.3) 26 (1.8) 25 (0.6)	30 (2.3) 24 (1.6) 27 (2.1) 27 (0.7)	
2 and over	71 (1.0)	24 (0.5)	26 (0.6)	23 (0.5)	20 (0.5)	25 (0.5)	25 (0.5)	24 (0.4)	25 (0.6)	27 (0.7)	
Non-Hispanic Asian:											
2 - 5	93* (4.6) 93* (3.2) 82 (4.1) 89 (2.0)	23* (1.6) 30 (1.9) 25 (2.2) 29 (1.0)	26* (2.6) 31 (1.7) 30 (2.1) 32 (1.0)	23* (1.5) 29 (2.1) 25 (2.4) 28 (0.9)	17* (2.0) 24 (3.9) 22 (4.1) 21 (0.9)	23* (2.4) 31 (2.6) 27 (1.8) 29 (1.0)	22* (2.0) 30 (2.0) 23 (2.2) 30 (1.3)	21* (1.6) 28 (2.0) 20 (2.4) 29 (1.6)	23* (2.6) 30 (1.9) 25 (2.1) 30 (1.4)	23* (2.7) 32 (2.9) 26 (2.4) 32 (1.5)	
2 and over	89 (1.5)	29 (1.0)	31 (0.9)	28 (0.9)	21 (0.9)	28 (0.9)	29 (1.3)	28 (1.5)	29 (1.3)	31 (1.4)	
Hispanic:											
2 - 5	79 (4.4) 77 (3.4) 78 (4.2) 69 (1.4)	21 (2.2) 24 (1.0) 28 (1.8) 23 (0.8)	25 (2.4) 27 (1.3) 32 (2.8) 27 (0.9)	19 (1.9) 22 (1.0) 26 (1.5) 22 (0.7)	16 (1.7) 20 (1.3) 23 (1.3) 18 (0.9)	23 (2.4) 25 (2.0) 29 (2.5) 24 (1.0)	23 (2.8) 25 (1.1) 29 (2.1) 24 (1.0)	21 (2.5) 23 (1.2) 28 (2.0) 24 (1.0)	24 (3.1) 26 (1.2) 29 (2.2) 25 (1.2)	27 (3.4) 26 (1.6) 30 (2.2) 25 (1.1)	
2 and over	72 (1.6)	24 (0.8)	27 (0.9)	22 (0.7)	19 (0.8)	25 (1.1)	25 (0.9)	24 (0.9)	25 (1.0)	26 (0.9)	

Table 18. Lunch¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic (continued)

Race/ethnicity and age (years)	Choles- terol % (SE)	Vitamin A (RAE) % (SE)	Beta- carotene % (SE)	Lycopene % (SE)	Thiamin % (SE)	Ribo- flavin % (SE)	Niacin % (SE)	Vitamin B6 % (SE)	Folate (DFE) % (SE)
Non-Hispanic White:									
2 - 5	23 (1.4)	19 (1.5)	29 (7.8)	31 (4.4)	21 (0.7)	21 (1.2)	27 (1.5)	21 (1.1)	19 (1.0)
6 - 11	23 (1.9)	20 (1.4)	29 (4.4)	43 (4.2)	23 (1.0)	23 (1.2)	27 (1.6)	22 (1.4)	20 (0.9)
12 - 19	25 (1.5)	21 (1.9)	29 (4.2)	24 (3.3)	25 (1.3)	23 (1.6)	27 (1.4)	23 (2.3)	22 (1.2)
20 and over	23 (0.8)	24 (1.0)	32 (1.8)	26 (2.1)	24 (0.4)	19 (0.8)	25 (0.7)	22 (0.8)	22 (0.7)
2 and over	23 (0.7)	23 (0.8)	32 (1.5)	27 (1.9)	24 (0.4)	20 (0.7)	26 (0.6)	22 (0.7)	22 (0.7)
Non-Hispanic Black:									
2 - 5	27 (1.6)	20 (2.4)	18 (5.1)	45 (8.4)	26 (1.7)	24 (1.9)	29 (1.6)	24 (1.7)	23 (1.5)
6 - 11	22 (1.7)	22 (1.4)	27 (3.1)	32 (4.8)	23 (1.4)	25 (1.4)	24 (1.1)	22 (1.2)	20 (2.0)
12 - 19	21 (1.8)	23 (1.6)	29 (2.8)	35 (4.8)	26 (2.0)	24 (1.5)	26 (1.7)	23 (1.4)	24 (2.1)
20 and over	22 (0.9)	21 (1.7)	29 (2.2)	27 (2.7)	24 (0.7)	21 (0.7)	25 (0.7)	23 (1.1)	23 (0.9)
2 and over	22 (0.9)	21 (1.5)	28 (2.0)	29 (1.8)	24 (0.5)	22 (0.6)	26 (0.6)	23 (0.9)	23 (0.5)
Non-Hispanic Asian:									
2 - 5	26* (3.0)	22* (5.6)	50* (18.8)	15* (4.7)	23* (2.2)	18* (1.6)	26* (2.9)	22* (2.5)	23* (2.1)
6 - 11	28 (2.3)	28 (2.8)	45 (6.7)	43* (5.0)	30 (2.2)	26 (2.1)	31 (1.5)	25 (1.6)	29 (1.8)
12 - 19	24 (2.1)	17 (2.0)	27 (4.6)	36 (7.1)	28 (3.1)	23 (2.9)	31 (2.1)	28 (2.0)	25 (1.8)
20 and over	31 (1.7)	29 (1.3)	36 (3.1)	42 (2.6)	30 (1.2)	25 (1.0)	32 (1.4)	29 (1.2)	31 (1.1)
2 and over	30 (1.5)	28 (0.8)	36 (2.9)	41 (2.1)	29 (1.3)	25 (1.1)	31 (1.3)	29 (1.1)	30 (1.0)
Hispanic:									
2 - 5	24 (4.2)	15 (1.7)	27 (5.2)	29 (8.1)	18 (1.8)	16 (1.8)	26 (2.4)	20 (1.9)	17 (2.1)
6 - 11	24 (2.6)	18 (1.5)	28 (3.6)	34 (4.5)	22 (1.0)	21 (1.0)	25 (0.9)	20 (1.1)	20 (1.0)
12 - 19	29 (3.5)	21 (2.6)	39 (4.8)	35 (3.3)	25 (2.1)	24 (2.2)	30 (2.3)	26 (2.2)	24 (1.9)
20 and over	23 (1.0)	21 (1.0)	30 (2.1)	30 (2.6)	24 (0.9)	19 (0.7)	25 (0.9)	22 (1.6)	23 (0.8)
2 and over	24 (1.1)	20 (1.0)	31 (1.8)	31 (2.2)	23 (0.8)	20 (0.7)	26 (0.8)	23 (1.3)	22 (0.8)

Table 18. Lunch¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic (*continued*)

Race/ethnicity										min E oha-								
and age	Cho	oline	Vitam	in B12	Vita	min C	Vitar	nin D	tocop	herol)	Vita	min K	Cal	cium	Phos	phorus	Magı	nesium
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:																		
2 - 5	22	(0.9)	18	(0.8)	26	(2.8)	16	(1.1)	28	(2.5)	28	(3.2)	21	(0.7)	24	(0.6)	23	(1.1)
6 - 11	23	(1.6)	21	(1.3)	25	(1.4)	19	(1.6)	27	(1.6)	27	(4.5)	26	(1.2)	26	(0.9)	25	(0.9)
12 - 19	24	(1.1)	22	(2.1)	18	(2.3)	20	(2.8)	22	(1.8)	27	(3.0)	25	(1.8)	25	(1.3)	22	(1.3)
20 and over	22	(0.6)	22	(1.2)	22	(0.8)	17	(0.8)	22	(0.8)	30	(1.3)	23	(0.5)	24	(0.5)	21	(0.5)
2 and over	22	(0.5)	22	(1.0)	22	(0.7)	17	(0.6)	23	(0.7)	30	(1.2)	23	(0.4)	24	(0.4)	21	(0.4)
Non-Hispanic Black:																		
2 - 5	27	(1.7)	23	(2.0)	24	(2.4)	20	(2.7)	28	(2.5)	21	(3.0)	25	(3.0)	28	(2.2)	26	(2.3)
6 - 11	24	(1.3)	24	(1.4)	26	(2.7)	25	(1.6)	23	(1.4)	22	(3.2)	28	(1.2)	27	(1.4)	25	(1.5)
12 - 19	23	(1.6)	25	(2.1)	24	(3.2)	20	(2.2)	26	(2.2)	26	(3.5)	28	(1.7)	28	(1.6)	27	(1.4)
20 and over	22	(0.8)	22	(1.6)	21	(1.3)	18	(1.9)	24	(0.8)	29	(1.9)	21	(0.8)	24	(0.6)	21	(0.6)
2 and over	22	(0.8)	22	(1.3)	22	(1.2)	19	(1.7)	24	(0.8)	28	(1.5)	23	(0.7)	25	(0.5)	22	(0.5)
Non-Hispanic Asian:																		
2 - 5	22*	(2.0)	15*	(2.8)	21*	(1.7)	12*	(2.0)	20*	(2.7)	24*	(3.9)	16*	(2.5)	21*	(2.3)	21*	(1.9)
6 - 11	29	(1.9)	22	(2.8)	24	(3.5)	20*	(2.0)	32	(2.5)	34	(3.5)	27	(2.6)	29	(1.9)	29	(1.7)
12 - 19	25	(2.3)	21	(2.1)	28	(3.7)	19	(5.5)	26	(1.9)	29	(4.2)	21	(3.5)	27	(2.3)	26	(2.2)
20 and over	29	(0.9)	27	(1.9)	28	(1.7)	25	(2.4)	30	(1.1)	38	(2.6)	24	(1.0)	29	(0.8)	27	(0.7)
2 and over	29	(0.9)	25	(1.6)	28	(1.5)	23	(1.7)	29	(0.9)	37	(2.3)	24	(1.0)	29	(0.8)	26	(0.7)
Hispanic:																		
2 - 5	21	(2.9)	16	(1.8)	20	(2.3)	11*	(1.5)	23	(2.7)	28	(4.1)	17	(1.4)	21	(2.0)	20	(2.0)
6 - 11	23	(1.7)	20	(1.5)	22	(3.3)	17	(1.8)	26	(1.8)	29	(3.5)	22	(1.6)	25	(1.2)	23	(1.4)
12 - 19	28	(2.5)	25	(3.2)	25	(2.5)	21	(3.8)	28	(2.0)	36	(3.5)	26	(2.2)	29	(2.3)	27	(2.1)
20 and over	22	(0.9)	21	(1.2)	21	(1.4)	16	(1.3)	23	(0.8)	27	(1.3)	20	(0.9)	24	(0.8)	21	(0.8)
2 and over	23	(1.0)	21	(1.1)	21	(1.1)	16	(1.4)	24	(0.8)	28	(1.2)	21	(0.9)	24	(0.9)	22	(0.8)

Table 18. Lunch¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic (continued)

D / 1 : :/																
Race/ethnicity	T.	on	7	inc	Co	nnor	Cala	nium	Doto	ssium	Soci	lium	Cof	feine	A 1a	cohol
and age (years)	и %	(SE)	Z .	(SE)	~ Co _j	pper (SE)	%	(SE)	Fota	(SE)	300 %	(SE)	Car.	(SE)	%	(SE)
(years)	70	(SE)	70	(SE)	70	(SE)	70	(SE)	70	(SE)	70	(SE)	70	(SE)	70	(SE)
Non-Hispanic White:																
2 - 5	20	(1.0)	22	(0.9)	22	(1.3)	27	(1.1)	24	(1.0)	29	(1.2)	27*	(11.6)		
6 - 11	22	(1.2)	25	(1.2)	25	(1.4)	29	(1.2)	25	(1.1)	30	(0.9)	16	(3.1)		
12 - 19	22	(1.1)	24	(1.1)	21	(1.4)	27	(1.3)	23	(1.2)	29	(1.4)	19	(3.0)		
20 and over	22	(0.5)	24	(0.6)	21	(0.6)	27	(0.5)	23	(0.5)	29	(0.5)	9	(0.9)	5	(0.9)
2 and over	22	(0.5)	24	(0.6)	21	(0.6)	27	(0.5)	23	(0.5)	29	(0.5)	9	(0.9)		
Non-Hispanic Black:																
2 - 5	22	(1.6)	25	(1.8)	27	(2.0)	32	(2.0)	28	(1.8)	33	(1.4)	23*	(8.8)		
6 - 11	21	(1.6)	24	(1.5)	24	(1.5)	25	(2.1)	27	(1.4)	26	(1.6)	13	(3.0)		
12 - 19	24	(1.8)	27	(1.6)	25	(1.6)	28	(2.0)	27	(1.5)	28	(1.7)	22	(4.0)		
20 and over	23	(0.8)	23	(0.7)	21	(0.8)	26	(0.7)	23	(0.7)	27	(0.6)	12	(0.9)	9*	(3.1)
2 and over	23	(0.5)	24	(0.6)	22	(0.6)	26	(0.6)	24	(0.6)	28	(0.6)	12	(0.9)		
Non-Hispanic Asian:																
2 - 5	20*	(2.4)	20*	(1.9)	22*	(1.8)	30*	(2.4)	21*	(1.8)	29*	(2.5)	9*	(3.7)		
6 - 11	28	(1.9)	28	(1.6)	30	(2.2)	33	(2.2)	30	(2.7)	33	(1.3)	17*	(4.9)		
12 - 19	25	(2.3)	27	(2.3)	27	(2.0)	30	(2.1)	29	(2.3)	29	(2.8)	14*	(3.8)		
20 and over	29	(1.3)	30	(1.1)	27	(0.8)	32	(1.1)	29	(0.8)	35	(1.5)	12	(1.3)	14*	(3.7)
2 and over	28	(1.2)	30	(1.1)	27	(0.7)	32	(1.0)	28	(0.7)	34	(1.5)	12	(1.3)		
Hispanic:																
2 - 5	17	(2.2)	20	(1.9)	22	(2.1)	27	(2.5)	21	(2.0)	29	(2.5)	9*	(4.4)		
6 - 11	20	(1.0)	24	(1.6)	24	(1.4)	27	(1.3)	24	(1.4)	27	(1.1)	22	(4.0)		
12 - 19	24	(1.9)	28	(2.5)	28	(2.3)	34	(3.0)	30	(2.4)	31	(2.3)	16	(2.5)		
20 and over	23	(0.8)	25	(1.0)	21	(0.8)	26	(0.8)	23	(0.9)	28	(0.9)	9	(0.6)	8	(1.6)
2 and over	22	(0.8)	25	(0.8)	22	(0.8)	28	(0.8)	24	(0.9)	28	(0.8)	10	(0.6)		

Symbol Legend

* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

Percent reporting: An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.76.

Nutrient ratios expressed as percentages: An estimated ratio between 25 and 75 percent is flagged when based on a sample size n^* of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and n^* is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of n^*p and n^* (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 2.76.

Footnotes

- ¹ Lunch includes eating occasions designated by the respondent as "brunch", "lunch" or the Spanish equivalent "comida." Please note these eating occasions include consumption of beverages including water.
- ² Percentages are estimated as a ratio of total nutrients from lunch for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: www.ars.usda.gov/nea/bhnrc/fsrg. See Table 2. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic.
- ³ The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item at an eating occasion designated as lunch.

Abbreviations

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2017-March 2020 Prepandemic

The 2017—March 2020 files represent a unique public-use data release from NHANES. The coronavirus disease 2019 (COVID-19) pandemic required suspension of data collection in March 2020. As a result, the partially completed NHANES 2019–2020 cycle was not nationally representative. Therefore, the 2019—March 2020 data were combined with the data from the 2017–2018 cycle to create the nationally representative 2017—March 2020 prepandemic data files.

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2017-March 2020 Prepandemic. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2017-2018 and 2019-2020 as applicable www.ars.usda.gov/nea/bhnrc/fsrg.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2022. Lunch: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, What We Eat in America, NHANES 2017-March 2020 Prepandemic.